# MATH327 – HOMEWORK SOLUTIONS HOMEWORK #4

Section 1.5: Problems 1, 2, 3, 5Section 2.1: Problems 1,4,7,9,10,14

Krzysztof Galicki

#### Problem 1.5.1.

- a) Yes, it is reflexive, antisymmetric, and transitive and therefore it is a partial order on  $\mathbb{R}$ . It is also a total order.
- b) No, as it is not reflexive. For example, it is not true that 2 < 2.
- c)  $(\mathbb{R}, \preceq)$ , where  $a \preceq b$  means  $a^2 \leq b^2$  is not a partial order as it is not antisymmetric. While  $a \prec b$  and  $b \prec a$  imply that  $a^2 = b^2$ , the latter does not mean that a = b.
- d) This relation is not antisymmetric. Consider (2,3) and (2,1). Clearly  $(2,3) \leq (2,1)$  and  $(2,1) \leq (2,3)$  but  $(2,3) \neq (2,1)$ .
- e) This is a partial order but not a total order. Consider, for example, (2,3) and (3,4). Neither  $(2,3) \prec (3,4)$ , nor  $(3,4) \prec (2,3)$ .
- f) This relation is not antisymmetric. Consider two words  $w_1 = cat$  and  $w_2 = pit$  (in this example the alphabet is the standard English alphabet). They have the same length so  $w_1 \leq w_2$  and  $w_2 \leq w_1$ . But they are not the same words.


## Problem 1.5.2.

- a) {1,10,100,1000,1001,101,1010,11,110,111}
- b) {1,11,111,110,10,101,1010,100,1001,1000}


#### **Problem 1.5.3.**

- a)  $a \prec b \prec c \prec d$  (other follow by transitivity).
- b)  $a \prec b$  and  $c \prec d$ .
- c)  $a \prec b$ ,  $a \prec d$ , and  $c \prec d$ .
- d)  $a \prec b$ ,  $a \prec c$ , and  $a \prec d$ .
- e)  $a \prec d \prec e, b \prec e,$  and  $b \prec c \prec f$  (other follow by transitivity).
- f)  $a \prec f$ ,  $g \prec f$ ,  $d \prec b \prec c$ ,  $d \prec h \prec i$ ,  $\prec c$ ,  $e \prec i$  (all the other relations follow by transitivity).


# Problem 1.5.5.



b)



(b') The Hesse diagram for  $(\mathcal{P}(\{a,b,c,d\}),\subseteq)$ 



(b) The Hesse diagram for  $(\{\{a\},\{a,b\},\{a,c\},\{c,d\},\{a,b,c\},\{a,b,c,d\}\},\subseteq)$ 

# Problem 2.1.1

- a) It is not a function as f(3) = 1 and f(3) = 3.
- b) It is not a function with domain  $\{1,2,3,4\}$  as f(3) is not defined.
- c) Yes, it is a function.
- d) It is not a function as f(1) = 1 and f(1) = 2.
- e) Yes, it is a function.

## Problem 2.1.4

- a) f(n) = 2n.
- b) f(n) = 1 + integer part of n/100.
- c)  $f(n) = n \mod 3 + 1.$
- d) f(n) = n.

# Problem 2.1.7

$$\{(1,-3),(3,1),(5,5),(2,5),(4,17)\}$$

The function is not 1-1 as f(5) = f(2).

#### Problem 2.1.9

- a) Write  $x^2 = |x|^2$  and the rest follows from the definition  $\sqrt{x^2} = \sqrt{|x|^2} = |x|$  (|x| is always non-negative).
- b) As both sides of this inequality are non-negative integers we can square it to get

$$|x+y| \le |x| + |y| \Leftrightarrow (x+y)^2 \le (|x|+|y|)^2 \Leftrightarrow x^2 + y^2 + 2xy \le x^2 + y^2 + 2|x||y| \Leftrightarrow xy \le |xy|.$$

The last inequality is obvious (you can verify by checking four cases of different sign of x and y).

### **Problem 2.1.10**

- a) It is not 1-1 as g(1) = g(-1). Neither it is onto as  $g(x) \ge 1$ .
- b) It is not 1-1 as g(1) = g(-1) but now it is clearly onto as any natural number can be written as |x| + 1. Simply take all positive integers and the zero for x.

**Problem 2.1.14** Write  $f(x) = (x+7)^2 - 100$ .

- a) 1-1 but not onto. f(n) is strictly increasing for  $n \ge -7$ , hence it must be 1-1. Also, there is no natural number n such that f(n) = -98. This would mean that  $f(n) = (n+7)^2 100 = -98$  or  $(n+7)^2 = 2$  which has no solutions in natural numbers.
- b) not 1-1 and not onto. To see this take f(-17) = f(-3) = 0, and observe that there is no integer such that f(m) = -98. This would mean that  $f(m) = (m+7)^2 100 = -98$  or  $(m+7)^2 = 2$  which has no integer solutions.
- c) not 1-1 but onto. Just draw this parabola in the XY-plane.